

FLASHES ON

Ejaculation pain

ATTIA ABDALLA ATTIA,MD.

**PROF. OF DERMATOLOGY & ANDROLOGY
FACULTY OF MEDICINE AL AZHAR UNV.**

Ejaculation pain

- Painful Ejaculation and post orgasmic pain are common conditions.
- which seriously affect:
 - sexual intercourse,
 - sexual satisfaction or
 - even the couple's relationship

- The intensity of pain ranges:
from simple discomfort
to non-specific pain
- Significantly impact the quality of life
- Its prevalence rate 1 – 4% amongst the general population.

Ejaculation pain

- It is a pelviperineal **pain** caused by ejaculation or orgasm.
- Mainly located in the penis
- Pain usually lasts < 5 minutes.

ejaculatory pain

COMMON CAUSES :

- Prostatitis
- Chronic pelvic pain syndrom
- LUTS-BPH
- Stones (Prostate: tubuloalveolar glands; SV.)

- Less important causes of painful ejaculation:

- *Ejaculatory duct obstruction*
- *Urethral stricture*
- *Psychogenic causes*
- *Prostate cancer*
- *Antidepressant therapy*
(tricyclic antidepressants, SSRIs, MAOIs)

TIME OF PAIN

- 1– AT EJACULATION
- Prostatitis
- Stones (in prostate &or seminal vesicle)
- Ejaculatory duct obstruction
- Urethral stricture

▪

TIME OF PAIN

2-AFTER EJACULATION

- Prostatitis
- Chronic pelvic pain syndrom
- LUTS-BPH

Prostatitis

- It is **defined** as inflammation of the prostate with an increased number of inflammatory cells within the prostate gland.
- *Tow Clacifications Of Prostatitis :*
 - Mears & Stamy (1978)
 - US- NIH (1995)

Prostatitis

(Mears & Stamy 1978)

- Acute Bacterial Prostatitis (ABP)**
- Chronic bacterial prostatitis (CBP)**
- Nonbacterial prostatitis**
- Prostatodynia.**

Prostatitis

(US- NIH 1995)

- Category I – ABP.
- Category II - Chronic bacterial prostatitis.
- Category III:(Ch. Pelvic Pain Syndrome)
 - a- Inflammatory CPPS. (ch abacterial p.)
 - b- Non Inflammatory CPPS . (prostatodynea)
- Category IIII : Asymptomatic Prostatitis

Prostatitis

PAIN in :

Genitourinary tract , the perineal area, penile tip, testicles, rectum, lower abdomen, and back.

• Urologic symptoms as :

frequency, urgency, dysuria, weak stream, nocturia, and postvoiding dribbling.

Chronic pelvic pain Syndrom

Is defined as: chronic pain,
pressure, or
discomfort

localized to: the pelvis,
perineum, or
genitalia of males

lasting more than 3 months

it is **not due to explainable causes:**

infection,
neoplasm, or
structural abnormality.

Lower Urinary Tract Symptoms (LUTS)

LUTS in men are categorised into:
voiding symptoms ,
storage symptoms or
post-micturition symptoms.

The most common cause of LUTS is benign
prostate hyperplasia (BPH), which obstructs
the bladder outlet.

LUTS

Voiding symptoms :

weak or intermittent urinary stream, straining, hesitancy, terminal dribbling and incomplete emptying.

Storage symptoms :

urgency, frequency, urgency incontinence and nocturia.

The major post-micturition symptoms :

post-micturition dribbling, which is common and bothersome.

THANK
YOU